

Table 4 Relative abundance of medium and large sized mammalian species recorded in the study area during dry and wet seasons

Species	Total number of individuals recorded		Relative abundance (%)	
	Dry	Wet	Dry	Wet
<i>C. abyssinicus</i> (colobus monkey)	95	70	9.84	9.27
<i>C. neglectus</i> (De brazza's monkey)	18	16	1.87	2.12
<i>C. aethiopsis</i> (vervet monkey)	148	107	15.34	14.17
<i>p. anubis</i> (olive baboon)	200	164	20.73	21.72
<i>H. cristata</i> (crested porcupine)	16	15	1.66	1.99
<i>L. fagani</i> (Bush hare)	16	14	1.66	1.85
<i>P. capensis</i> (rock hyrax)	28	16	2.90	2.12
<i>O. afer</i> (aardvark)	11	21	1.14	2.78
<i>T. scriptus</i> (common bushbuck)	27	13	2.80	1.72
<i>R. arundinum</i> (Common reed buck)	15	11	1.55	1.46
<i>S. grimmia</i> (Grey Duiker)	15	11	1.55	1.46
<i>P. larvatus</i> (bush pig)	24	28	2.49	3.71
<i>P. africanus</i> (warthog)	18	13	1.87	1.72
<i>S. caffer</i> (African buffalo)	213	183	22.07	24.24
<i>A. buselaphus</i> (Tora harte beast)	18	14	1.87	1.85
<i>H. hynae</i> (striped hyena)	13	7	1.35	0.93
<i>F. servestris</i> (African wild cat)	24	11	2.49	1.46
<i>C. mesamolas</i> (blackback jackal)	21	9	2.18	1.19
<i>I. albicauda</i> (white tailed mongoose)	10	13	1.04	1.72
<i>M. capensis</i> (honey badger)	14	8	1.45	1.06
<i>C. civetta</i> (African civet)	12	8	1.24	1.06
<i>P. pardus</i> (leopard)	5	1	0.52	0.13
<i>P. leo</i> (lion)	4	2	0.41	0.26
Total	965	755	100%	100%